

			temps de suivis	publié par	titre article	nb de cas	nb implant	impl perdu	Impl rajoutés apr	CRS av rabout	taux de succès impl CRS	CRS proth	date de l'étude	auteur	
State of the art implant placement and restoration through guided techniques															
Fast&fixed			2004-2007 : 3 ans	European Journal for Dental Implantologists	Implantation in the atrophied Jaw	55	292	6			97%		2008 F Kistler S Kistler G Bayer	Martin Marcus DDS, Russell Baer DDS, Nelson Lo DDS Chicago Illinois	
Fast&fixed			5	Ste internet		95	575	5					Weiss	Ulm Allemagne	
fast and fix			5			68	272	6			99%	100%	Delorme Carotte Geog Bayer	Landsberg Allemagne	
implants full arch			10 ans	Clin Oral Implants Res	Ten-year survival rates of fixed prostheses on four or six implants ad modum Bränemark in full edentulism.	156		4					Bränemark PI, Svensson B, van Steenberghe D, Pär-Olov Östman, DDS; * Mats Hellman, DDS; *Lars Sennerby, DDS,	Göteborg Suède	
				Clin implant dentistry and related research	Direct Implant Loading in the Edentulous Maxilla Using a Bone Density		123	1			99%	100%	2005 PhD*	Suède	
all on for six	maxillaire		5				51	204			98.36				Suisse
all on for six	mandibule						73	292			99.73%				
quattro	maxillaire			Clin Oral Implants Res									M, Villa C, Malo P		
fast and fix	mandibule														
	mandibule		févr-13	Département de chirurgie buccale, Faculté de médecine dentaire, Université de Milan, Paolo S. Institut des sciences biomédicales, en Italie. Clin Oral Implants Res	Faculté de médecine	226	904			96,8	98,5	21 997 Chiapasco M			
4 impl interforaminal	Mandibule						82	328			96,1		Chiapasco M, Gatti C.		
all on for six				Clin Dental Impl Res		44	176	62	96,7	98,2	100	2003 Malo P, Ranger , Nobre			
all on for six			1 an	Clin Implant Dentistry Related Res	A Pilot Study of Complete Edentulous Rehabilitation with Immediate Function Using a New Implant Design: Case Series	53	189			98,9		2005 Malo p	Portugal	All-on-Four™ immediate-function concept with Bränemark System implants to completely edentulous mandibles: a retrospective clinical study	
				Clin Implant Dentistry Related Res	Immediate loading in oral implants. Present situation.									Conclusions: The results	
					Immediate loading of immediate mandibular implants with a full-arch fixed prosthesis: a preliminary study.	10	60			100%	100%	2009 Periarrocha M, Boronat A, Garcia B.	Espagne		
4 impl interforaminal et 2 postérieurs			1997 à 2002	Medicina oral, Pathología oral, y cirugía bucal											
4 impl interforaminal et 2 postérieurs			12 mois	Journal of oral and maxillofacial surgery											
4 impl mand 4 ou 6 maxillaire				International oral & maxillofacial implants	Immediate loading of 190 endosteal dental implants: a prospective observational study of 40 patient treatments with up to 2-year data.	40	190			100%	100%	Nikellis I, Levi A, Nicolopoulos C.	Grèce		
4 impl interforaminal MCI amovible sur barre	mand prothèse amovible sur barre		36 à 96 mois	Clinical Implant dentistry and related research	Implant-retained mandibular overdentures with immediate loading: a 3 to 8-year prospective study on 328 implants.	1119	328	7	2	88,2	97,6 99%	100%	2003 Chiapasco M, Gatti C.	Italie	The results of this limited investigation suggest that patients who are partially or completely edentulous may be immediately restored with implants and fixed provisional restorations, provided that the dental implants are adequately stable immediately after their surgical placement. This alternative therapeutic approach did not appear to affect the up-to-2-year survival of the implants in this patient population.